

## Lung Rescue Unit

<u>Guidelines</u> for initiation and management of VV ECMO

(non lung-transplant/non cardiac-surgery/non trauma patients)

## **Inclusion Criteria**

- 1. Murray Score ≥ 3 (cesar.lshtm.ac.uk/murrayscorecalculator.htm)
  - a. Enter # of quadrants with infiltrates on CXR
  - b. Enter  $PaO_2/FiO_2$  in mmHg on 100% for 20 minutes
  - c. Enter PEEP value (cm H<sub>2</sub>O)
  - d. Enter compliance (ml/cm H<sub>2</sub>O) [Tidal volume/(PIP-PEEP)]
- 2. Hypercapnia (CO<sub>2</sub>>60) with pH <7.25, or inability to adequately ventilate with Pplat  $\leq$  30 cm H<sub>2</sub>O
- 3. On ventilator  $\leq$  10 days
- 4.  $\leq$  65 years of age
- 5. Patient with a *reversal* form of ARDS (e.g., infectious, trauma, post-operative)
- 6. Bedside physician clinical discretion

## **Exclusion criteria (Relative)**

- 1. > 65 years of age
- 2. > 10 days on ventilator
- 3. Requirement for home  $O_2$  therapy for severe lung disease
- 4. Severe neurological insult / injury
- 5. Inability to tolerate anticoagulation
- 6. Terminal disease with low 1-year survival rates
- 7. Severe underlying liver disease
- 8. Acute hepatic failure
- 9. Ongoing /uncontrolled hemorrhage
- 10. Jehovah's Witness (unwilling to receive blood products)
- 11. Acquired Immune Deficiency Syndrome (AIDS) related-cause of respiratory failure
- 12. Bedside physician clinical discretion