



UNIVERSITY of MARYLAND
UPPER CHESAPEAKE HEALTH

American College of Surgeons Commission on Cancer CP3R Data
Standards 4.4 and 4.5

CP ³ R Measures		Target	Goal	2013	2014	2015	2016	2017	2018
				UCMC	UCMC	UCMC	UCMC	UCMC	UCMC
Breast	Radiation therapy is administered within 1 year(365 days) of diagnosis for women under age 70 receiving breast conserving surgery for breast cancer. [BCS/RT]	90%	100%	97%	100%	100%	97%	* 98%	100%
	Combination chemotherapy is considered or administered within 4 months (120 days) of diagnosis for women under 70 with AJCC T1c NO MO, or Stage II or III hormone receptor negative breast CA. [MAC]	90%	100%	100%	100%	100%	100%	100%	100%
	Tamoxifen or third generation aromatase inhibitor is considered or administered within 1 yr (365 days) of diagnosis for women with AJCC T1c N0 MO, or Stage II or III hormone receptor positive breast CA. [HT]	90%	100%	100%	100%	100%	100%	100%	100%
	Needle core or FNA biopsy is performed prior to surgical treatment of breast cancer. [NB]	80%	90%	100%	95%	97%	100%	** 97%	100%
	Radiation therapy is considered or administered within 1 year of diagnosis for women with AJCC Stage III breast cancer undergoing modified radical mastectomy with four or more positive regional lymph nodes. [MASRT]	90%	100%	100%	100%	NA	100%	100%	100%
	Breast conservation therapy is performed for women with AJCC Stge 0, I, or II breast cancer [BCS]	Not defined	50%	61%	88%	92%	90%	*** 91%	*** 87%

American College of Surgeons Commission on Cancer CP3R Data
Standards 4.4 and 4.5

CP ³ R Measures		Target	Goal	2014	2016	2017	2018	
				UCMC	UCMC	UCMC	UCMC	
Colon	Adjuvant chemotherapy is considered or administered within 4 months (120 days) of diagnosis for patients under the age of 80 with AJCC Stage III (lymph node positive) colon cancer. [ACT]	90%	100%	100%	100%	100%	100%	
	At least 12 regional lymph nodes are removed and pathologically examined for resected colon cancer. [12RLN]	85%	100%	97%	96%	100%	* 90%	* 100%
Rectal	Preoperative chemo and radiation are administered for clinical AJCC T3N0, T4N0, or Stage III; or postoperative chemo and radiation are administered within 180 days of diagnosis for clinical AJCC T1-2 N0 with pathologic AJCC T3N0, T4N0, or Stage III; or treatment is recommended; for patients under the age of 80 receiving resection for rectal cancer. [RECTCT]	85%	100%	75%	100%	100%	100%	
Gastric	At least 15 regional lymph nodes are removed and pathologically examined for resected gastric cancer. [G15RLN]	80%		NA	100%	NA	NA	
NSCLC	At least 10 regional lymph nodes are removed and pathologically examined for AJCC stages IA, IB, IIA, and IIB resected NSCLC. [10RLN]	Not defined		42%	80%	72%	** 89%	** 68%
	Systemic chemotherapy is administered within 4 months to day preoperatively or day of surgery to 6 months postoperatively, or is considered for surgically resected cases with pathologic lymph node-positive (pN1) and (pN2) NSCLC. [LCT]	85%		100%	100%	100%	100%	
	Surgery is not the first treatment for cN2 M0 NSCLC cases. [LN0Surg]	85%			100%	100%	100%	



UNIVERSITY of MARYLAND
UPPER CHESAPEAKE HEALTH

American College of Surgeons Commission on Cancer CP3R Data

Standards 4.4 and 4.5

CP ³ R Measures		Target	Goal				2016	2017	2018
				UCMC	UCMC	UCMC	UCMC	UCMC	UCMC
CERVICAL	Radiation therapy completed within 60 days of initiation of radiation among women diagnosed with any stage of cervical cancer. [CERRT]	Not defined				100%	100%	100%	100%
	Chemotherapy administered to cervical cancer patients who received radiation for stages IB2-IV cancer (Group1) or with positive pelvic nodes, positive surgical margin, and/or positive parametrium (Group 2). [CERCT]	Not defined				50%	100%	100%	*
	Use of brachytherapy in patients treated with primary radiation with curative intent in any stage of cervical cancer. [CBRRT]	Not defined					0%	100%	**
OVARIAN	Salpingo-oophorectomy with omentectomy, debulking; cytoreductive surgery, or pelvic exenteration in Stages I-IIIc Ovarian cancer. [OVSAL]	Not defined				100%	NA	NA	***
ENDOMETRIAL	Chemotherapy and/or radiation administered to patients with Stage IIIc or IV Endometrial cancer. [ENDCTR]	Not defined				100%	100%	NA	NA
	Endoscopic, laproscopic, or robotic surgery performed for all Endometrial cancer (excluding sarcoma and lymphoma), for all stages except stage IV. [ENDLRC]	Not defined				NA	50%	100%	100%

American College of Surgeons Commission on Cancer CP3R Data

Standards 4.4 and 4.5

CP ³ R Measures		Target	Goal				2016	2017	2018
				UCMC	UCMC	UCMC	UCMC	UCMC	UCMC
BLADDER	At least 2 lymph nodes are removed in bladder cancer patients under 80 undergoing partial or radical cystectomy. (BL2RLN)	Not defined						NA	NA
	Radical or partial cystectomy; or Tri-modality therapy (Local tumor destruction/excision with chemotherapy and radiation) for clinical T234N0M0 patients, first treatment within 90 days of diagnosis (BLCSTRI)	Not defined						*	67%
	Neo-adjuvant or adjuvant chemotherapy offered or administered for patients with muscle invasive cancer undergoing radical cystectomy (BLCT)	Not defined						NA	NA